

**\*\*FULL TITLE\*\***

*ASP Conference Series, Vol. \*\*VOLUME\*\*, \*\*YEAR OF PUBLICATION\*\**

**\*\*NAMES OF EDITORS\*\***

## The formation of extreme mass ratio binary stars: a tribute to Eduardo Delgado Donate

C. J. Clarke

*Institute of Astronomy, Cambridge, CB3 0HA, U.K.*

**Abstract.** Eduardo Delgado was due to have presented a poster at this meeting on his latest results on the formation of extreme mass ratio binaries. Tragically, Eduardo was among those killed in a hiking accident in Tenerife earlier this year. As his PhD supervisor, and as a longstanding collaborator, the organisers of this meeting kindly invited me to incorporate a report on his most recent work into a more general tribute to his life and work. I will reflect on Eduardo's scientific career, the problems that motivated him and his achievements, focusing particularly on a problem which had intrigued us both for several years and on which Eduardo was making important progress at the time of his death. Finally, I will mention the personal qualities that Eduardo brought to his work and the acute sense of loss that is shared by all those - friends and collaborators - who were privileged to know him.

### 1. Cambridge days: predictions and puzzles in multiple star formation

Eduardo came to Cambridge from his native Tenerife in 2000, in order to work on hydrodynamic (SPH) star formation simulations under my supervision. He was funded by an E.U. PhD studentship as part of the 'Young Stellar Cluster' Network. Anyone who has been involved in these Networks will know that it is not always easy to find students of suitable calibre to fill these posts, given their restrictive nationality and residence requirements, and can therefore imagine my delight at receiving such a strong application from Eduardo. He went on to fulfil my high expectations, quickly developing into a fine numericist and astrophysicist. (I should add that I was often struck by the fact that Eduardo's written English was far more eloquent than anything that his British contemporaries could manage - although I'm afraid that this says something not only about Eduardo but also about the British educational system!)

Eduardo arrived in Cambridge when Matthew Bate was still a postdoc there and Matthew was a great help to Eduardo in helping him to master his SPH code. This was an exciting time for Eduardo to be working in numerical star formation, as this was the time that Matthew was producing the first turbulent fragmentation calculations which have generated so much interest (and controversy!) ever since. It was clearly going to be a challenge for Eduardo to do anything in this field which was going to have a distinct impact. I think the collection of well cited papers that Eduardo produced during his thesis shows that he met this challenge most successfully.

Eduardo became known in the field for his series of turbulent fragmentation calculations which explored the production of multiple star systems and brown dwarfs in small N ensembles. My own vote for the part of his thesis that will be




Figure 1. Examples of some of the system architectures of complex multiple star systems formed in the hydrodynamical simulations of Delgado et al 2004. The numbers refer to separations in A.U..

his biggest contribution to the field would go to his calculations which were the first to make predictions about the properties of higher order multiple systems (see some of the complex system architectures generated in Figure 1). He was able to do this because rather than focusing his computational effort on system scale (as Matthew Bate did with his 50 solar mass calculation), Eduardo followed small (5 solar mass) clouds and - in compensation - was able to follow the calculation for much longer. Thus Eduardo's simulations could follow even wide systems to the point of dynamical stability. Evidently this is important, not only because wide binaries are numerous in nature (half of all binaries with solar type primaries are wider than 30 A.U.) but because in this way one can start to make pronouncements about higher order multiples (which necessarily involve wide, as well as close, components).

Thus Eduardo entered uncharted waters in terms of simulations (and indeed his paper on this, Delgado et al 2004 is still the only one on this topic in existence). Moreover, the characterisation of multiple star systems is far from complete observationally, as it is a laborious task to collate complete statistics on components with a wide dynamic range of separations (e.g. Tokovinin et al 2006). Indeed, the motivation for such observational studies has always been undermined by the fact that - until Eduardo's work - there were no theoretical predictions with which to compare the data.

This brings me to one of Eduardo's most important characteristics as a scientist, that in addition to being technically accomplished as a numericist, he had a strong interest in - and respect for - observations. One would never hear Eduardo implying that observations much be wrong if they contradicted his simulations - or that observers should simply go and 'look harder'. He understood, better than most, that it's actually the areas of disagreement that illuminate your calculations. And, moreover, he understood that the range of diagnostic information that is contained in multiple star statistics would ultimately provide a very stringent - arguably the most stringent - test of star formation theories.

Thus Eduardo became intrigued by an obvious area of discrepancy between observations and all star formation simulations. In the simulations, all the binaries were within a factor of two of being of unit mass ratio (with the exception of a few weakly bound very low mass 'outliers' at large separations). Moreover, this result extended to every level of the hierarchy (i.e. in a triple, a nearly equal mass binary would be bound to a third star with mass nearly equal to the binary). Evidently nature does not work like that! It is true that there are categories of binaries that favour more nearly equal masses, arguably OB stars (Garcia & Mermilliod 2001), very low mass binaries (Bouy et al 2003) and short period spectroscopic binaries among solar type stars (Mazeh et al 1992). However, for the bulk of solar type binaries, the median mass ratio is  $\sim 0.4$  (Duquennoy & Mayor 1991), a result which - given that it is based partly on pairs from the visual binary literature which are almost certainly selectively incomplete at low mass ratios - must, if anything, *underestimate* the incidence of pairs with very disparate masses.

Early in 2003, Eduardo started writing a paper with me, alas never to be published, entitled 'On the problem of forming extreme mass ratio binaries.'. I told him to start by writing a few pages setting up the problem (which he did) and then, by the time he'd written that, his simulations would have run to the point where he could unveil the solution (or at any rate *a* solution) in the concluding sections. But the latter proved to be impossible. None of the fixes to initial conditions which he attempted (such as adjusting the power spectrum of the initial velocity field or changing the geometry) were ultimately successful. He would excitedly tell me that yes, he'd managed to make a binary with  $q$  of 0.1, only to return a week later to tell me that, no, it had now evolved to  $q$  of 0.7.... Thus it turned out that the problem was not the initial creation of such systems, but the effect of continued accretion onto the protobinary, which drove up the mass ratio.

It has been known since at least the work of Artymowicz 1981 (and note that Eduardo went on to spend two years in Stockholm working with Pawel Artymowicz as a postdoc, although never on this problem) that accretion of gas whose specific angular momentum exceeds that of the binary leads to preferential accretion onto the secondary. This commonsense result, which was apparent in these early ballistic calculations and was backed up by the subsequent SPH simulations of Bate & Bonnell 1997 and Bate 2000, occurs simply because the secondary's Roche lobe is further from the system's centre of mass, and is thus more likely to intercept high angular momentum material. Since in the case of any plausible binary formation scenario, higher angular momentum material will fall in later, the binary mass ratio,  $q$ , should therefore rise with time.

## 2. Numerical controversy: latest results

While Eduardo and I abandoned this question, discouraged, (and Eduardo took up a position in Stockholm where he worked mainly on codes for planet formation), there meanwhile appeared the paper of Ochi, Sugimoto & Hanawa 2005. This claimed, using a grid based code, that actually material of high specific angular momentum is preferentially accreted on to the binary *primary* and thus that the mass ratio should *fall*. This study did not contradict the widely acknowledged result that the flow preferentially enters the secondary's Roche lobe, but argued instead that, after half an orbit around the secondary, it crosses via the L1 point into the primary's Roche lobe and is then accreted. Ochi et al conjectured that the failure of SPH calculations to demonstrate this behaviour stemmed from the excessively viscous nature of SPH, which might cause the SPH particles to spiral in sufficiently, in half an orbit, so that they could avoid the L1 region. In the same spirit, proponents of SPH countered that Ochi et al's flows were too warm (with sound speed of  $0.25\times$  the orbital velocity of the binary) and that this caused artificial acceleration of the flow through the L1 point. Clearly these issues had to be investigated further, using both Lagrangian and Eulerian codes. At the time of his death, Eduardo was close to completing a suite of high resolution SPH calculations which were beginning to illuminate the problem and whose results I summarise below.

His repetition, at higher resolution, of the SPH calculation of Bonnell & Bate 1997 (which modeled accretion onto a protobinary of a flow with fixed specific angular momentum ) demonstrated that the rate of increase of  $q$  is indeed somewhat resolution dependent and is over-estimated at low resolution (about  $5 \times 10^4$  particles in the discs) compared with the highest resolution (a hundred times more particles, which has achieved numerical convergence) by about a factor two. This can be traced to the fact that at low resolution, particles in the outer part of the primary's Roche lobe are not behaving in a fully fluid-like manner and, through precession of their elliptical orbits, are able to tip back through the L1 point into the secondary's Roche lobe. Nevertheless,  $q$  *increases* (for this value of specific angular momentum of the flow) *at all resolutions*, in contrast to the results of Ochi et al.

Eduardo found that at high resolution, the shocks are well delineated (see left hand panel of Figure 2) and, indeed, that the Jacobi constant (i.e. the Bernoulli function in the co-rotating frame) is well conserved along streamlines except in shocks. This result goes against the hypothesis of Ochi et al that the SPH results are driven by artificial viscosity in the secondary's disc flow, since under these circumstances the action of artificial viscosity should cause changes in the Jacobi constant, even away from shocks. Crucially, however, the shock geometry is quite different in Eduardo's simulations compared with those of Ochi et al. In the SPH calculation, loosely bound material in the secondary's Roche lobe encounters a shock with material flowing in from the direction of the primary (see dense structure to the East of the secondary in left hand panel of Figure 2) and it is dissipation in this shock which prevents material flowing through L1 and into the primary's Roche lobe. This shock is simply not there in Ochi et al's calculation, so that material entering via L2 retains enough energy to smoothly transit through L1.




Figure 2. Density and flow structure in the co-rotating frame around an accreting proto-binary, contrasting a cold three dimensional simulation (left) and a warm two dimensional simulation (right). The secondary is to the left. Note the pronounced shock to the East of the secondary in the left hand panel. Unpublished simulations by E. Delgado

Thus while both simulations are apparently self-consistent, *given the location of the shocks*, one has to enquire why the shock morphology is so different in the two cases. One possibility is that - although the SPH and Eulerian simulations are matched in terms of specific angular momentum of the accretion flow and mass ratio of the binary - there are two potentially important differences. Firstly, the Eulerian calculation is much warmer (about a factor 25 in temperature). Secondly, it is strictly two dimensional whereas the SPH calculation - although highly flattened near the binary - is fully three dimensional and introduces particles on the surface of a sphere at large radius. An indication that one, or both, of these effects may be important is provided by one of Eduardo's simulations which mimicked as closely as possible the precise conditions of Ochi et al, being both 2D and warm. The right hand panel of Figure 2 demonstrates that the shock has in this case moved away from the secondary's Roche lobe, in qualitative agreement with the flow morphology of Ochi et al. Although *quantitative* agreement between the SPH and Eulerian codes is not achieved even here, the *sign* of the change is encouraging. This shift in flow morphology is reflected in a change in the relative accretion rates on to the primary and secondary. Although  $\dot{q}$  still increases in these warm 2D SPH calculations,  $\dot{q}$  is reduced by an order of magnitude compared with the cold 3D calculation.

This was the situation at the time of Eduardo's death and there is evidently still much for his collaborators to do. We now suspect that gas temperature

and/or flow dimensionality are important determinants of the shock morphology, and we must now discover which of these two effects is the critical one. One then has to judge which set of conditions is more realistic. Ultimately, this will tell us whether the old ‘simple’ idea is correct - i.e. that the flow simply enters the secondary’s Roche lobe and remains therein - or whether the more complex outcome of Ochi et al - involving flow between Roche lobes - is closer to the truth.

All this has implications for the extreme mass ratio binary problem which Eduardo had been trying to solve for years. If the result is closer to that of Ochi et al, then this could solve the problem at a stroke. If the results instead support the old SPH results, then the puzzle remains (although if low resolution simulations have over-estimated the *magnitude* of the growth of  $q$ , then binaries may remain at low  $q$ , even if the sign of  $\dot{q}$  is positive). Otherwise, one has to think harder about how to prevent the infall of gas with high specific angular momentum onto protobinaries. Although it is tempting to invoke feedback for this, it is not obvious why this should be so important in the low mass binaries which we are largely trying to explain.

### 3. Eduardo the person

Having talked about the problems that motivated Eduardo as a scientist, it leads me to how much can one discern of Eduardo the person in all this? An unexpected death leads one to analyse what one saw of a person in the daily contacts of studentship and collaboration. Eduardo was always calm, always gentle, always attentive. He was never arrogant, never distorted the problem at hand, or his relations with those around him, by the obtrusion of his own ego. He commanded the respect of all those who came into contact with him through this obvious intellectual and personal integrity. He was a joy to work with.

I see great continuity between those characteristics as a collaborator and the personal testimonies from his friends, many of which are movingly collected on his blogspot (<http://eduardo-delgado.blogspot.com/>) I was aware that Eduardo was very popular with his peers in Cambridge and that he belonged to a big crowd of friends of all nationalities whose joyful social life never got in the way of Eduardo’s scientific activities. But these messages - and the evidence of the conversations I have had with so many - speak of a very special warmth of affection and regard. Unfortunately I can only dimly decipher the majority in Spanish but they clearly tell the same story as those I can read (in Italian and English). So many speak of the gentleness of his eyes, the warmth of his friendship, his calmness (one Italian friend exclaimed ‘You even played football with calm’ (incredible!)). He was, to put it simply, much loved.

All this is very particular to Eduardo, but I should end on a note that perhaps has relevance to all those gathered for this meeting and to astronomers everywhere. In the last months of his life, Eduardo was particularly happy. A former Cambridge colleague, who stayed with him on an observing run at Christmas time, told me ‘You know, Edu was just *so* happy when I last saw him’. He had everything he dreamed of, living in the island he loved, starting new projects in his new postdoc with Casiana Munoz, also continuing his independent research (he felt he was so close to cracking the extreme mass ratio binary

problem). We're perhaps not all in one of those special moments in life when everything seems to be working, but I reckon that if any one of us suffered the same awful and random misfortune as Eduardo, then they would be able to say of us that we'd spent our lives doing what we loved. This is a rare privilege. So, in keeping with the celebratory mood of this conference, let us recall that, while mourning Eduardo no less, we can also celebrate a life spent - like ours - in doing something we believe to be intrinsically valuable. To do this in a spirit of calm and integrity was Eduardo's gift. We miss him immensely.



Figure 3. Edu emanating his characteristic calm

## References

- Artymowicz, P. 1983 *Acta Astron.* 33,223
- Delgado-Donate, E. J., Clarke, C. J., Bate, M. R. & Hodgkin, S. T. 2004, *MNRAS* 351,617
- Duquennoy, A. & Mayor, M. 1991, *A & A* 248,485
- Bate, M. R. 2000, *MNRAS* 314,33
- Bate, M. R & Bonnell, I. A. 1997, *MNRAS* 285,33
- Bouy, H., Brandner, W., Martin, E., Delfosse, X., Allard, F. & Basri, G. 2003, *AJ* 126,1526

Mazeh T., Goldberg D., Duquennoy A., Mayor M. 1992, ApJ, 401, 265

Garcia, B. & Mermilliod, J. 2001, A & A 368,122

Ochi, Y., Sugimoto, K. & Hanawa, T. 2005, ApJ 623,922

Tokovinin, A., Thomas, S., Sterzik, M. & Udry, S. 2006, A & A 450,681